

Children's Motrin/Advil (IBUPROFEN)		Tylenol (ACETAMINOPHEN)	
<ul style="list-style-type: none"> ➔ *DO NOT GIVE TO CHILDREN UNDER 6 MONTHS OF AGE* ➔ Motrin is to be given every 6 – 8 hours ➔ DO NOT exceed 4 doses in a 24-hour period. ➔ ALWAYS use the measuring device provided. ➔ While taking this medication encourage fluid intake. ➔ Give with food. ➔ *INFANT'S Motrin is <i>half</i> of the amounts listed below 		<ul style="list-style-type: none"> ➔ *DO NOT GIVE TO CHILDREN UNDER 2 MONTHS OF AGE * ➔ Tylenol is to be given every 4 – 6 hours ➔ DO NOT exceed 5 doses in a 24-hour period. ➔ ALWAYS use the measuring device provided. 	
Dosing Chart for Children's Motrin *not Infant's Motrin		Dosing chart for Children's Tylenol	
12 -17 lbs	2 – 3 mL	12 -17 lbs	2 .5- 3.5 mL
18 - 23 lbs	3 – 5 mL	18 - 23 lbs	3.5 – 5 mL
24 - 35 lbs	5 – 8 mL	24 - 35 lbs	6 – 7.5 mL
36 – 47 lbs	8 – 11 mL	36 – 47 lbs	8 – 10 mL
48 – 59 lbs	12– 13 mL	48 – 59 lbs	10– 12.5 mL
60 – 71 lbs	13.5 – 16 mL	60 – 71 lbs	12.5 – 15 mL
72 – 95 lbs	16 – 20 mL	72 – 95 lbs	15 -20 mL

These medications are weight-based, and dosing is based on the child's weight in kilograms (converted to pounds). The doses are presented in ranges that are considered safe even without knowing any specifics about the child. Additionally, the dose is typically rounded to an amount that is easy to administer while still being within a safe range.

In the event of a dosage error, please call Poison Control: 1-800-222-1222

DIPHENHYDRAMINE (BENADRYL) DOSING TABLE					
Child's Weight (pounds) / Type of Benadryl	20-24	25-37	38-49	50-99	100+
LIQUID (12.5mg/5mL)	4ml	5ml	7.5ml	10ml	_____
LIQUID (12.5mg/1 teaspoon)	¾ tsp	1tsp	1.5 tsp	2 tsp	_____
CHEWABLE (12.5mg)	_____	1 tablet	1.5 tablets	2 tablets	4 tablets
TABLETS (25mg)	_____	½ tablet	½ tablet	1 tablet	2 tablets
CAPSULES (25mg)	_____	_____	_____	1 capsule	2 capsules