# PUCILLO FAMILY PRACTICE P.A. WELLNESS CENTER

Family Medicine & Bariatrics


# PFP March 2022: COVID updates, COVID Booster, Colorectal health







Dr. Ron Pucillo, M.D.



Dr. Brian Tong, M.D.

Why is March the most popular month to use a trampoline? It is spring-time Which March flowers grow on faces?


Tulips (2 lips)

### February 2022 newsletter:


- COVID updates
- Vaccine and booster updates
- March colorectal cancer month

# **COVID updates**

- New Omicron variants
  - o Omicron BA.2 is a descendant of original Omicron BA.1.
    - it is 30% more transmissible than the original Omicron strain
    - becoming the dominant strain in the U.S. as of March 31st
    - Vaccines and booster protection Based on early evidence, vaccines and booster is 40% effective protection for infection and 80-90% for hospitalization



Reinfection - WHO indicated those with previous BA.1 infection provides strong protection against reinfection with BA.2




- XE variant
  - initially found in UK. it is a recombinant mutation of BA.1 and BA.2. causing it to be more transmissible
  - more information to come

# **Vaccine and Booster update**



- On March 29, 2022, FDA approved Moderna and Pfizer booster vaccine (4th shot) to individuals >50 years old with chronic medical conditions
  - Chronic medical conditions include cancer, CKD, chronic liver disease, chronic lung disease, cystic fibrosis, dementia, diabetes, organ transplant
  - o booster may be given at least 4 months after the 1st booster.
  - Below is a graph that measures the effectiveness of a 4th dose.
     (NEJM, below the dashed line indicates + diagnosis of COVID cases)



Vaccine and Booster schedule


Vaccine	0 monti	1 mont	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month
Pfizer-BioNTech (ages 5-11 years)	1" Dose		3 <sup>rd</sup> Dose (At least 4 weeks after 2 <sup>rd</sup> dose)							
Pfizer-BioNTech (ages 12 years and older)	1" Dose	2 <sup>nd</sup> Dose (3 weeks after 1 <sup>st</sup> dose)	3 <sup>rd</sup> Dose (At least 4 weeks after 2 <sup>rd</sup> dose)			poster Dose <sup>1</sup> least 3 months after 3	3 <sup>rd</sup> dose)			Booster Dose <sup>3</sup> e footnote)
Moderna (ages 18 years and older)	1" Dose	2 <sup>rd</sup> Dose (4 weeks after 1 <sup>st</sup> dose)	3rd Dose (At least 4 weeks after 2rd dose)			Booster Dose <sup>1</sup> (at least 3 months a	after 3 <sup>rd</sup> dose)			2 <sup>nd</sup> Booster Dose <sup>3</sup> (See footnote)
Janssen (ages 18 years and older)	1 <sup>st</sup> Dose	an mRNA CO	al) Dose <sup>2</sup> using VID-19 Vaccine ks after 1 <sup>st</sup> dose)	Booster Dose <sup>1</sup> (at least 2 months after additional dos	e)			2 <sup>nd</sup> Booster Dose (See footnote)		

# **Colorectal Health**

### March is Colorectal Health Month

### Colon anatomy and function

- colon is divided into cecum, ascending colon, transverse colon, descending colon, sigmoid colon and rectum
- it's functions to absorb nutrients, electrolytes and water




#### Colon health

 Reduces risk of colon cancer healthy weight, physically active, calcium and vitamin D, quit smoking, eating vegetables, fruits, whole grains.



## Colon cancer screening

- Colonoscopy every 3, 5, 10 years depending on the findings
- Flexible sigmoidoscopy every 5 years
- CT colonography every 5 years
- Stool test
  - o FOBT yearly
  - Cologuard every 3 years



We thank you for allowing us to be part of your family in 2022. Please feel free to contact us if you have any questions.

Sincerely, Staff at Pucillo Family Practice